

Fleischforschung und Entwicklung

## Die Produktion nicht kastrierter Eber – Herausforderungen und Chancen

Von Ronald E. Klont, Esra Kurt, Lourens Heres und Bert Urlings

*Schlüsselwörter:* nicht kastrierte Eber | Ebergeruch | Verringerung | Prävention | Erkennung

### Literatur

1. BABOL, J., SQUIRES, E.J., und K. LUNDSTRÖM (1996): Investigation of factors responsible for the development of boar taint. *Food Research International* 28, 573-581.
2. BANON, S., E. COSTA, M.D. GIL und M.-D. GARRIDO (2003): A comparative study of boar taint in cooked and dry-cured meat. *Meat Sci.* 63, 381–88.
3. BONNEAU, M., KEMPSTER, A. J., CLAUS, R., CLAUDI-MAGNUSSEN, C., DIESTRE, A., TORNBERG, E., WALSTRA, P., CHEVILLON, P., WEILER, U. und G.L. COOK (2000): An international study on the importance of androstenone and skatole for boar taint: I. Presentation of the programme and measurement of boar taint compounds with different analytical procedures. *Meat Sci.* 54, 251–259.
4. BONNEAU, M., P. WALSTRA, C. CLAUDI-MAGNUSSEN, A.J. KEMPSTER, E. TORNBERG, K. FISCHER, A. DIESTRE, F. SIRET, P. CHEVILLON, R. CLAUS, G. DIJKSTERHUIS, P. PUNTER, K. R. MATTHEWS, H. AGERHEM, M. P. BÉAGUE, M. A. OLIVER, M. GISPERT, U. WEILER, G. VON SETH, H. LEASK, M. FONT I FURNOLS, D. B. HOMER und G. L. COOK (2000): An international study on the importance of androstenone and skatole for boar taint: IV. Simulation studies on consumer dissatisfaction with entire male pork and the effect of sorting carcasses on the slaughter line, main conclusions and recommendations. *Meat Sci.* 54, 285–295.
5. CLAUS, R., A. FISCHER und B. VOGELBACHER (1985): Konzentrationen des Ebergeruchsteroids im Schlachtkörper des Ebers und daraus hergestellter Fleischzeugnisse. *Fleischwirtsch.* 65 (3), 375–377.
6. CLAUS, R., WEILER, U., und A. HERZOG (1994): Physiological aspects of androstenone and skatole formation in the boar: a review with experimental data. *Meat sci.* 38, 289–305.
7. DIJKSTERHUIS, G.B., B. ENGEL, P. WALSTRA, M. FONT I FURNOLS, H. AGERHEM, K. FISCHER, M.A. OLIVER, C. CLAUDI-MAGNUSSEN, F. SIRET, M.P. BEAGUE, D.B. HOMER und M. BONNEAU (2000): An international study on the importance of androstenone and skatole for boar taint: II. Sensory evaluation by trained panels in seven European countries. *Meat Sci.* 54, 261–269.
8. DUNSHEA, F.R., C. COLANTONI, K. HOWARD, I. MCCAULEY, P. JACKSON, K.A. LONG, S. LOPATICKI, E.A. NUGENT, J.A. SIMONS, J. WALKER und D.P. HENNESEY (2001): Vaccination of boars with GnRH vaccine (Improvac) and eliminates boar taint and increases growth performance. *Journal of Animal Sci.* 79, 2524–2535.
9. FONT I FURNOLS M., J. GONZÁLEZ, M. GISPERT, M.A. OLIVER, M. HORTÓS, J. PÉREZ, P. SUÁREZ, L. und GUERRERO (2009): Sensory characterization of meat from pigs vaccinated against gonadotropin releasing factor compared to meat from surgically castrated, entire male and female pigs. *Meat Sci.* 83, 438–442.
10. GERRITZEN M.A., M. KLUIVERS-POODT, H. G. M. REIMERT, V. HINDLE und E. LAMBOOIJ (2008): Castration of piglets under CO<sub>2</sub>-gas anaesthesia *Animal*, 2:11, pp 1666–1673.

11. GERRITZEN M.A., M. KLUIVERS-POODT, H. G. M. REIMERT, V. HINDLE und E. LAMBOOIJ (2009): Response to the Letter to the Editor on the surgical castration of piglets. *Animal*, 3:11, pp 1476–1477.
12. HAUGEN, J-E. (2003): Rapid detection of boar taint with chemical sensor technology. The challenge of gas-sensor based methods. European Association of Animal Production Working Group on Production and Utilisation of Meat from Entire Males, 13-14 November, Dublin, Ireland.
13. HUBER-EICHER, B. und P. SPRING (2008): Attitudes of Swiss consumers towards meat from entire or immunocastrated boars: A representative survey. *Research in Veterinary Sci.* 85, 626–627.
14. JOHNSON, L.A., RATH, D. VASQUES, J.M., MAXWELL und J.R. DOBRINSKY (2005): Preselection of sex of offspring in swine for production: current status of the process and its application. *Theriogenology*, 63, 615–624.
15. KLUIVERS-POODT, M. H. HOPSTER und H.A.M SPOOLDER (2007): Castration under anaesthesia and/or analgesia in commercial pig production. Report 85, Animal Science Group Lelystad, Wageningen-UR, The Netherlands.
16. LEE, G.J. A.L. ARCHIBALD, A.S. LAW, S. LLOYD, J. WOOD und C.S. HALEY (2005): Detection of quantitative trait loci for androstenonen, skatole and boar taint between large white and Meishan pigs. *Animal Genetics* 36, 14–22.
17. LUNDE K., B. EGELANDSDAL, J. CHOINSKI, M. MIELNIK, A. FLATTEN und E. KUBBEROD (2008): Marinating as a technology to shift sensory thresholds in ready-to-eat entire male pork meat. *Meat Sci.* 80, 1264–1272.
18. MALMFORS, B. und I. HANSON (1974): Incidence of boar taint in Swedish Landrace and Yorkshire boars. *Livestock Production Sci.*, 1, 411–420.
19. MALMFORS, B. und K. LUNDSTRÖM (1983): Consumer reactions to boar taint, a review. *Livestock production Sci.* 10, 187–196
20. MATTHEWS, K. R., HOMER, D. B., PUNTER, P., BEAGUE, M.P., GISPERT, M., KEMPSTER, A.J., AGERHEM, H., CLAUDI-MAGNUSSEN, C., FISCHER, K., SIRET, F., LEASK, H., FONT I FURNOLS, M. und M. BONNEAU (2000): An international study on the importance of androstenonen and skatole for boar taint: III. Consumer survey in seven European countries. *Meat Sci.* 54, 271–283.
21. MCCAULEY, I., D.P. HENNESEY, V. BOGHOSSION, L. SALI, L. SALVATORE, J. REYNOLDS und R. MAWSON (1997): Effects of cooking and processing pork on the perception of boar taint. In Bonneau, M, K. Lundstrom, & B. Malmfors (editors), *Boar taint in entire male pigs*. E.A.A.P Publication no. 92, Wageningen Pers.
22. MOE, M., T. MEUWISSEN, S. LIEN, C. BENDIXEN, X. WANG, L.N. CONLEY, I. BERGET, H. TAJET und E. GRINDFLEK (2007): Gene expression profiles in testis of pigs with extreme high and low levels of androstenone. *BMC Genomics* 8, doi:10.1186/1471-2164-8-405.
23. MOE, M., L. S. LIEN, T. AASMUNDSTAD, T. H.E. MEUWISSEN, M.H.S. HANSEN, C. BENDIXEN und E. GRINDFLEK (2009) Association between SNPs within candidate genes and compounds related to boar taint and reproduction *BMC Genetics* 2009, 10:32.
24. MORTENSEN, A.B. und S.E. SORENSEN (1984): Relationship between boar taint and skatole determined with a new analysis method. 30th European meeting of Meat Research Workers, Bristol, United Kingdom.
25. MOTTRAM, D.S., J.D. WOOD und R.L.S. PATTERSON (1982): Comparison of bulls and castrates for bacon production. 3. Composition and eating quality of bacon. *Animal Production* 35, 75-80.
26. MÜHLBAUER, I. W. OTTEN, W. LÜPPING, A. PALZER, S. ZÖLS, S. ELICKER, M. RITZMANN und K. HERINTITZL (2009): Investigation about CO<sub>2</sub> gas anaesthesia

- as an alternative option for piglet castration without anaesthesia. *Der Praktische Tierarzt* 90 (5) 460-464.
27. PAULY P., J.V. SPRING, O. DOHERTY, S. AMPUERO KRAGTEN und G. BEE (2008): Performances, meat quality and boar taint of castrates and entire male pigs fed a standard and a raw potato starch-enriched diet. *Animal* 2:11, 1707–1715.
  28. QUINTANILLA, R., O. DEMEURE, J. P. BIDANEL, D. MILAN, N. IANNUCELLI, Y. AMIGUES, J. GRUAND, C. RENARD, C. CHEVALET und M. BONNEAU (2003): Detection of quantitative trait loci for fat androstenone levels in pigs. *J. Anim. Sci.* 81, 385–394.
  29. RATH, D. (2002): Low dose insemination in the sow. A review. *Reproduction in Domestic Animals*, 37, 201-205.
  30. STOLZENBACH, S., G. LINDAHL, K. LUNDSTRÖM, G. CHEN und D.V. BYRNE (2009): Perceptual masking of boar taint in Swedish fermented sausages. *Meat Sci.* 81, 580–588.
  31. SQUIRES, E.J. (2006): Possibilities for selection against boar taint. *Acta Veterinaria Scandinavica*, 48 (Suppl 1), S8.
  32. TAJET, H., O. ANDRESEN und T.H.E. MEUWISSEN (2006): Estimation of genetic parameters of boar taint; skatole and androstenone and their correlations with sexual maturation. *Acta Veterinaria Scandinavica*, 48 (Suppl 1), S9.
  33. TER BEEK, V. (2007) Sex sorting technology could save time and money. *Pig Progress*, 23 (10), 22–23.
  34. VARONA, L., O. VIDAL, R. QUINTANILLA, M. GIL, A. SÁNCHEZ, J. M. FOLCH, M. HORTOS, M. A. RIUS, M. AMILLS und J. L. NOGUERA (2005): Bayesian analysis of quantitative trait loci for boar taint in a Landrace outbred population. *Journal of Animal Science* 83, 301–307.
  35. WALSTRA, P. (1974): Fattening of young boars: quantification of negative and positive aspects. *Livestock Production Science*, 1, 187–196.
  36. WALSTRA, P., C. CLAUDI-MAGNUSSEN, P. CHEVILLON, G. VON-SETH, A. DIESTRE, K.R. MATTHEWS, D.B. HOMER, and M. BONNEAU (1999): An international study on the importance of androstenone and skatole for boar taint: levels of androstenonen and skatole by country and season. *Livestock Production Science*, 62, 15-28.
  37. WEILER, U. M. FONT I FURNOLS, K. FISCHER, H. KEMMER, M.A. OLIVER, M. GISPERTT, A. DOBROWOLSKI und R. CLAUS (2000): Influence of differences in sensitivity of Spanish and German consumers to perceive androstenonen on the acceptance of boar meat differing in skatole and androstenone concentrations. *Meat Sci.* 54, 297-304.
  38. WILLEKE, H. (1993): Possibilities of breeding for low 5 $\alpha$ -androstenone content in pigs. *Pig News and Information*, 14, 31-33.
  39. WILLIAMS, L. D., PEARSON, A. M. und N. B. WEBB (1963): Incidence of sex odor in boars, sows, barrows, and gilts. *Journal of Animal Sci.* 22, 116–168.
  40. WRIGHT A.J., M. WHITING und A. TAYLOR (2009): Letter to the Editor on the surgical castration of piglets. *Animal*. 3 (11), 1474–1475.
  41. XUE, J. L., DIAL, G.D., HOLTON, E.E., VICKERS, Z., SQUIRES, E.J., LOU, Y., GOTBOUT, D. and N. MOREL (1996): Breed differences in boar taint: relationship between tissue levels of boar taint compounds and sensory analysis. *Journal of Animal Sci.* 74, 2170–2177.

### **Anschrift der Verfasser**

Dr. Ronald E. Klont, Dr. Esra Kurt, Dr. Manager R&D Food Safety Lourens Heres und Prof. Dr. Bert Urlings, Vion Food Group, 5280 AA Boxtel, Niederlande  
(R. E. Klont ist korrespondierender Autor, ronald.klont@vionfood.com)